

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
4 October 2001 (04.10.2001)

PCT

(10) International Publication Number  
**WO 01/72132 A1**

(51) International Patent Classification<sup>7</sup>: **A01N 63/00**,  
65/00, A61K 48/00, G01N 33/00, A01K 67/00, 67/027,  
67/033, C12N 15/00

Ludwig Institute for Cancer Research, P.O. Box 240, S-171  
77 Stockholm (SE).

(21) International Application Number: PCT/US01/09855

(74) Agent: **EVANS, Joseph, D.**; Evenson, McKeown, Ed-  
wards & Lenahan, P.L.L.C., 1200 G Street, N.W., Suite  
700, Washington, DC 20005 (US).

(22) International Filing Date: 28 March 2001 (28.03.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/192,507 28 March 2000 (28.03.2000) US

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,  
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,  
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,  
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,  
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,  
TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(71) Applicant: **LUDWIG INSTITUTE FOR CANCER RE-  
SEARCH** [US/US]; 605 Third Avenue, New York, NY  
10158 (US).

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,  
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors: **ERIKSSON, Ulf**; Ludwig Institute for Can-  
cer Research, P.O. Box 240, S-171 77 Stockholm (SE). **LI,  
Xuri**; Ludwig Institute for Cancer Research, P.O. Box 240,  
S-171 77 Stockholm (SE). **PONTEN, Annica**; Ludwig In-  
stitute for Cancer Research, P.O. Box 240, S-171 77 Stock-  
holm (SE). **AASE, Karin**; Ludwig Institute for Cancer Re-  
search, P.O. Box 240, S-171 77 Stockholm (SE). **LI, Hong**;

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: NON-HUMAN TRANSGENIC ANIMALS EXPRESSING PLATELET-DERIVED GROWTH FACTOR C (PDGF-C)  
AND USES THEREOF

(57) Abstract: Non-human transgenic animals overexpressing PDGF-C and cells thereof have been created. The transgenic animals contain a nucleotide sequence that encodes for platelet derived growth factor C (PDGF-C) or an analog thereof. These animals are useful for studying disease states characterized by overexpression of PDGF-C, as well as useful for evaluating therapies intended to treat such diseases.

WO 01/72132 A1

BEST AVAILABLE COPY

NON-HUMAN TRANSGENIC ANIMALS EXPRESSING PLATELET-DERIVED  
GROWTH FACTOR C (PDGF-C) AND USES THEREOF

This application claims the priority of co-pending U.S. provisional application Serial No. 60/192,507 filed March 28, 2000 in the name of Ulf ERIKSSON et al.

5        This invention relates to non-human transgenic animals which overexpress PDGF-C and cells thereof that are useful for research on the effects that overexpression of PDGF-C has on an organism, and particularly for assaying substances which inhibit PDGF-C activity.

10

Background of the Invention

Platelet-derived growth factors (PDGFs) are important in the growth, survival and function of connective tissue cells, fibroblasts, myofibroblasts and glial cells (Heldin et al., Growth Factor, 1993 8 245-252). In adults, PDGFs stimulate wound healing (Robson et al., Lancet, 1992 339 23-25). Structurally, PDGF isoforms are disulfide-bonded dimers of homologous A- and B-polypeptide chains, arranged as homodimers (PDGF-AA and PDGF-BB) or as a heterodimer (PDGF-AB).

20

PDGF isoforms exert their effects on target cells by binding to two structurally related receptor tyrosine kinases (RTKs). The alpha-receptor (PDGFR-alpha) binds both the A- and B-chains of PDGF, whereas the beta-receptor (PDGFR-beta) binds only the B-chain. These two receptors are expressed by many cell lines grown *in vitro*, and are mainly expressed *in vivo* by mesenchymal cells. The PDGFs exert their effects *in vivo* in a paracrine mode since they often are expressed in epithelial (PDGF-A) or endothelial (PDGF-B) cells in close apposition to the PDGFR-expressing mesenchyme. In tumor cells and in cell lines grown in

25

30

*vitro*, coexpression of the PDGFs and the PDGFRs generates autocrine loops which are important for cellular transformation (Betsholtz et al., Cell, 1984 39 447-57; Keating et al., J. R. Coll Surg Edinb., 1990 35 172-4).

5 Overexpression of the PDGFs has been observed in several pathological conditions, including malignancies, arteriosclerosis, and fibroproliferative diseases (reviewed in Heldin et al., The Molecular and Cellular Biology of Wound Repair, New York: Plenum Press, 1996, 249-273).

10 The importance of the PDGFs as regulators of cell proliferation and survival is well illustrated by recent gene targeting studies in mice that have shown distinct physiological roles for the PDGFs and their receptors despite the overlapping ligand specificities of the PDGFRs.

15 Homozygous null mutations for either of the two PDGF ligands or the receptors are lethal. Approximately 50% of the homozygous PDGF-A deficient mice have an early lethal phenotype before embryonic day E10. The surviving animals have a complex postnatal phenotype with lung emphysema due  
20 to improper alveolar septum formation because of a lack of alveolar myofibroblasts (Boström et al., Cell, 1996 85 863-873). The PDGF-A deficient mice also have a dermal phenotype characterized by thin dermis, misshapen hair follicles and thin hair (Karlsson et al., Development, 1999  
25 126 2611-2). PDGF-A is also required for normal development of oligodendrocytes and subsequent myelination of the central nervous system (Fruttiger et al., Development, 1999 126 457-67).

30 The phenotype of PDGFR-alpha deficient mice is more severe with incomplete cephalic closure, impaired neural crest development, cardiovascular defects, skeletal defects and edemas, leading to embryonic death around E8-16

(Soriano *et al.*, Development, 1997 124 2691-70). The PDGF-B and PDGFR-beta deficient mice develop similar phenotypes that are characterized by renal, hematological and cardiovascular abnormalities and death at E17-19 (Levéen *et al.*, Genes Dev., 1994 8 1875-1887; Soriano *et al.*, Genes Dev., 1994 8 1888-96; Lindahl *et al.*, Science, 1997 277 242-5; Lindahl, Development, 1998 125 3313-2). The renal and cardiovascular defects are due, at least in part, to the lack of proper recruitment of mural cells (vascular smooth muscle cells, pericytes or mesangial cells) to blood vessels (Levéen *et al.*, Genes Dev., 1994 8 1875-1887; Lindahl *et al.*, Science, 1997 277 242-5; Lindahl *et al.*, Development, 1998 125 3313-2).

The PDGFs are members of the Platelet Derived Growth Factors/Vascular Endothelial Growth Factors (PDGF/VEGF) family of growth factors which presently consists of nine different members. The members of the PDGF/VEGF family are all characterized by the presence of eight conserved cysteine residues. In their active, physiological state, these proteins are dimers formed by disulfide bonding, by both inter- and intramolecular bonds, at the eight cysteine residues.

Besides PDGF-A and PDGF-B, the members of this family include VEGF and five proteins that are closely related to VEGF, and a new factor related to the PDGFs, designated PDGF-C. The five proteins closely related to VEGF are: VEGF-B, described in International Patent Application PCT/US96/02957 (WO 96/26736) which corresponds to U.S. Patent 5,928,939 and in U.S. Patents 5,840,693 and 5,607,918 by Ludwig Institute for Cancer Research and The University of Helsinki; VEGF-C or VEGF2, described in Joukov *et al.*, EMBO J., 1996 15 290-298 and Lee *et al.*,

Proc. Natl. Acad. Sci. USA, 1996 93 1988-1992, and U.S. Patents 5,932,540, 5,935,540 and 6,040,157 by Human Genome Sciences, Inc; VEGF-D, described in International Patent Application No. PCT/US97/14696 (WO 98/07832), and Achen et al., Proc. Natl. Acad. Sci. USA, 1998 95 548-553; the placenta growth factor (PlGF), described in Maglione et al., Proc. Natl. Acad. Sci. USA, 1991 88 9267-9271; and VEGF3, described in International Patent Application No. PCT/US95/07283 (WO 96/39421) by Human Genome Sciences, Inc.

Each VEGF family member has between 30% and 45% amino acid sequence identity with VEGF. Functional characteristics of the VEGF and the VEGF-related proteins include varying degrees of mitogenicity for endothelial cells, induction of vascular permeability and angiogenic and lymphangiogenic properties.

Similarity between two proteins is determined by comparing the amino acid sequence and conserved amino acid substitutions of one of the proteins to the sequence of the second protein, whereas identity is determined without including the conserved amino acid substitutions.

VEGF is a homodimeric glycoprotein that has been isolated from several sources. Alternative mRNA splicing of a single VEGF gene gives rise to five isoforms of VEGF. VEGF shows highly specific mitogenic activity for endothelial cells. VEGF has important regulatory functions in the formation of new blood vessels during embryonic vasculogenesis and in angiogenesis during adult life (Carmeliet et al., Nature, 1996 380 435-439; Ferrara et al., Nature, 1996 380 439-442; reviewed in Ferrara and Davis-Smyth, Endocrine Rev., 1997 18 4-25). The significance of the role played by VEGF has been demonstrated in studies showing that inactivation of a

single VEGF allele results in embryonic lethality due to failed development of the vasculature (Carmeliet et al., Nature, 1996 380 435-439; Ferrara et al., Nature, 1996 380 439-442). The isolation and properties of VEGF have been  
5 reviewed; see Ferrara et al., J. Cellular Biochem., 1991 47 211-218 and Connolly, J. Cellular Biochem., 1991 47 219-223.

In addition VEGF has strong chemoattractant activity towards monocytes, can induce the plasminogen activator and  
10 the plasminogen activator inhibitor in endothelial cells, and can also induce microvascular permeability. Because of the latter activity, it is sometimes referred to as vascular permeability factor (VPF). VEGF is also chemotactic for certain hematopoietic cells. Recent  
15 literature indicates that VEGF blocks maturation of dendritic cells and thereby reduces the effectiveness of the immune response to tumors (many tumors secrete VEGF) (Gabrilovich et al., Blood, 1998 92 4150-4166 and Gabrilovich et al., Clinical Cancer Research, 1999 5 2963-  
20 2970).

VEGF-B has similar angiogenic and other properties to those of VEGF, but differs from VEGF in its distribution and expression in tissues. In particular, VEGF-B is very strongly expressed in heart and weakly in lung, whereas the  
25 reverse is the case for VEGF. This suggests that VEGF and VEGF-B, despite the fact that they are co-expressed in many tissues, may have functional differences.

VEGF-B was isolated using a yeast co-hybrid interaction trap screening technique by screening for  
30 cellular proteins which might interact with cellular retinoid acid-binding protein type I (CRABP-I). Its isolation and characteristics are described in detail in

PCT/US96/02957 (WO 96/26736), in U.S. Patents 5,840,693 and 5,607,918 by Ludwig Institute for Cancer Research and The University of Helsinki and in Olofsson et al., Proc. Natl. Acad. Sci. USA, 1996 93 2576-2581.

5 VEGF-C was isolated from conditioned media of the PC-3 prostate adenocarcinoma cell line (CRL1435) by screening for ability of the medium to produce tyrosine phosphorylation of the endothelial cell-specific receptor tyrosine kinase VEGFR-3 (Flt4), using cells transfected to  
10 express VEGFR-3. VEGF-C was purified using affinity chromatography with recombinant VEGFR-3, and was cloned from a PC-3 cDNA library. Its isolation and characteristics are described in detail in Joukov et al., EMBO J., 1996 15 290-298.

15 VEGF-D was isolated from a human breast cDNA library, commercially available from Clontech, by screening with an expressed sequence tag obtained from a human cDNA library designated "Soares Breast 3NbHBst" as a hybridization probe (Achen et al., Proc. Natl. Acad. Sci. USA, 1998 95 548-  
20 553). Its isolation and characteristics are described in detail in International Patent Application No. PCT/US97/14696 (WO98/07832).

The VEGF-D gene is broadly expressed in the adult human, but is certainly not ubiquitously expressed. VEGF-D  
25 is strongly expressed in heart, lung and skeletal muscle. Intermediate levels of VEGF-D are expressed in spleen, ovary, small intestine and colon, and a lower expression occurs in kidney, pancreas, thymus, prostate and testis. No VEGF-D mRNA was detected in RNA from brain, placenta,  
30 liver or peripheral blood leukocytes.

PlGF was isolated from a term placenta cDNA library. Its isolation and characteristics are described in detail in Maglione et al., Proc. Natl. Acad. Sci. USA, 1991 88

9267-9271. Presently its biological function is not well understood.

5 VEGF3 was isolated from a cDNA library derived from colon tissue. VEGF3 is stated to have about 36% identity and 66% similarity to VEGF. The method of isolation of the gene encoding VEGF3 is unclear and no characterization of the biological activity is disclosed.

10 As with the PDGFs, the VEGF family members act primarily by binding to receptor tyrosine kinases. Five endothelial cell-specific receptor tyrosine kinases have been identified, namely VEGFR-1 (Flt-1), VEGFR-2 (KDR/Flk-1), VEGFR-3 (Flt4), Tie and Tek/Tie-2. All of these have the intrinsic tyrosine kinase activity which is necessary for signal transduction. The essential, specific  
15 role in vasculogenesis and angiogenesis of VEGFR-1, VEGFR-2, VEGFR-3, Tie and Tek/Tie-2 has been demonstrated by targeted mutations inactivating these receptors in mouse embryos.

20 Most recently, an additional member of the PDGF/VEGF family of growth factors was identified, which is called PDGF-C. PDGF-C is described in International Patent Application PCT/US99/22668, filed September 30, 1999, in co-pending U.S. Application Serial no. 09/410,349, filed September 30, 1999, as well as in co-pending U.S.  
25 Provisional Application Serial No. 60/192,507, filed March 28, 2000, which are all specifically incorporated herein by reference.

30 PDGF-C has a two-domain structure not previously recognized within this family of growth factors, an N-terminal C1r/C1s/embryonic sea urchin protein Uegf/bone morphogenetic protein 1 (CUB) domain, and a C-terminal PDGF/VEGF homology domain (P/VHD). The structure of the P/VHD in PDGF-C shows a low overall sequence identity with



other PDGF/VEGF homology domains, although the eight invariant cysteine residues involved in inter- and intra-molecular disulfide bond formation are present. The cysteine spacing in the central, most conserved region of this domain is different from other PDGF/VEGF domains, with an insertion of three amino acid residues. Despite the fact that the insertion occurs close to the loop 2 region which has been proposed to be involved in receptor binding, it was shown that this domain of PDGF-CC dimers binds PDGFR-alpha with an affinity almost identical to homodimers of PDGF-A or -B chains. In addition, four extra cysteine residues are present in this domain. Full length and truncated PDGF-CC dimers were found not to bind to VEGFR-1, -2 or -3, or to PDGFR-beta.

PDGF-C requires proteolytic removal of the N-terminal CUB domain for receptor binding and activation of the receptor. This indicates that the CUB domains are likely to sterically block the receptor binding epitopes of the unprocessed dimer. The *in vitro* and *in vivo* proteolytically processed proteins are devoid of N-terminal portions corresponding to more than 14-16 kDa as determined from SDS-PAGE analysis that is consistent with a loss of the 110 amino acid long CUB domain and a variable length portion of the hinge region between the CUB and core domains.

PDGF-C is not proteolytically processed during secretion in transfected COS cells indicating that proteolytic removal of the CUB domain occurs extracellularly, and not during secretion. This is in contrast to PDGF-A and -B (Östman et al., J. Cell. Biol., 1992 118 509-519) which appear to be processed intracellularly by furin-like endoproteases (Nakayama et al., Biochem J., 1997 327 625-635).

*In situ* localization studies demonstrate expression of PDGF-C in certain epithelial structures, and PDGFR-alpha in adjacent mesenchyme, indicating the potential of paracrine signaling in the developing embryo. PDGF-C expression  
5 seems particularly abundant at sites of ongoing ductal morphogenesis, indicating a role of the factor in connective tissue remodeling at these sites. The expression pattern is distinct from that of PDGF-A or PDGF-B indicating that the three growth factors have different  
10 roles despite their similar PDGFR-alpha binding and signaling activities. This is illustrated by the mouse embryonic kidney, in which PDGF-C is expressed in early aggregates of metanephric mesenchyme undergoing epithelial conversion, whereas PDGF-A is expressed in more mature  
15 tubular structures, and PDGF-B by vascular endothelial cells. PDGFR-alpha is expressed in the mesenchyme of the kidney cortex, adjacent to the sites of PDGF-C expression, indicating that this mesenchyme may be targeted specifically by PDGF-C. Indeed, PDGFR-alpha -/- mouse  
20 embryos show an extensive loss of the cortical mesenchyme adjacent to sites of PDGF-C expression, not seen in PDGF-A -/- mice or in PDGF-A/B -/- mice, indicating that PDGF-C has an essential role in the development of kidney mesenchyme.

25 Northern blots show PDGF-C mRNA in a variety of human tissues, including heart, liver, kidney, pancreas and ovary.

Transgenic manipulation can result in overexpression of a protein, making transgenic animal models useful tools  
30 to study the functions and physiological activities of proteins. A variety of such animal models have been produced for this purpose. One technique for producing transgenic animals involves the process of microinjection

of a foreign DNA or transgene into the pronuclei of a fertilized egg. The introduced DNA appears to integrate randomly into the chromosome. Another technique for producing transgenic animals involves modifying an embryonic stem cell to overexpress a transgene.

#### Summary of the Invention

In one of its aspects, the invention involves overexpression of PDGF-C or an analog thereof, or a functional activated fragment of PDGF-C or an analog thereof in non-human transgenic animals. In particular, the transgenic animals of the invention are useful in both understanding the effects of overexpressing PDGF-C and as a research tool for developing compounds that will inhibit the effects caused by overexpression of PDGF-C, such as development of hypertrophy and fibrosis in various organs, such as the heart.

The term "analog" or "functional analog" refers to a modified form of PDGF-C in which at least one amino acid substitution has been made such that the analog or functional analog retains substantially the same biological activity as the unmodified PDGF-C *in vivo* and/or *in vitro*.

There are many "biological activities of PDGF-C" that can be readily tested by methods known in the art. For example, the polypeptide or the encoded polypeptide from a polynucleotide has the ability to stimulate one or more of proliferation, differentiation, motility, survival or vascular permeability of cells expressing a PDGF-C receptor including, but not limited to, vascular endothelial cells, lymphatic endothelial cells, connective tissue cells (such as fibroblasts), myofibroblasts and glial cells.

Preferably the polypeptide or the encoded polypeptide from a polynucleotide has the ability to stimulate wound healing. PDGF-C can also have antagonistic effects on cells.

5 As used herein, the term "PDGF-C" collectively refers to the polypeptides of SEQ ID NO:1 or SEQ ID NO:2, and fragments or analogs thereof which have the biological activities of PDGF-C as defined above, and to a polynucleotide which encodes PDGF-C, or a fragment or  
10 analog thereof having the biological activities of PDGF-C.

In another aspect, the invention provides a method for producing a transgenic, non-human animal overexpressing PDGF-C or an analog thereof, or a functional activated fragment of PDGF-C or an analog thereof. One method  
15 comprises introducing a transgenic DNA into the pronuclei of a fertilized egg of a non-human animal. The transgenic DNA is operably linked to a promoter. The egg is then implanted into a pseudopregnant non-human animal and allowed to develop into a transgenic animal. Examples of  
20 promoters that can be used include, but are not limited to, the alpha-myosin heavy chain promoter which gives cardiac myocyte-specific expression, the keratin K14 promoter which gives basal keratinocyte-specific expression and the insulin promoter which gives pancreatic beta cell-specific  
25 expression.

An alternative method of producing such a transgenic non-human animal is to modify an embryonic stem cell to overexpress PDGF-C or an analog thereof, or a functional activated fragment of PDGF-C or an analog thereof. This  
30 method comprises introducing a transgenic DNA into embryonic stem cells of a non-human animal. This cell is introduced into an animal embryo at a stage when the cell is capable of integrating, for example, at the blastocyte

stage. The embryo is allowed to develop into a transgenic animal.

Subsequent to the production of a first generation of transgenic animals, a further alternative method is to mate  
5 a transgenic animal containing the sequences described here or made available by this invention with a second animal.

For example, a transgenic mouse according to the present invention can be mated with a wild-type mouse to  
10 produce 50% wild type and 50% heterozygous transgenic mice. The heterozygous transgenic mice offspring overexpress PDGF-C to the same degree as the parental generation.

Further, a transgenic mouse according to the present invention can be mated with a mouse transgenic for a trait  
15 other than PDGF-C overexpression. For example, the tsT A58 transgenic mouse (Immortomouse) that expresses the SV40 T antigen could be used, see Jat et. al., PNAS Vol. 88 pages 5096-5100 (1991). Cell lines isolated from a resultant mouse would be conditionally immortal. These cell lines  
20 would multiply when grown at 33°C. If the temperature was raised to 39°C, the cells would differentiate, allowing growth of cell lines such as cardiac myocytes. Cell lines could be isolated for the purpose of use in an assay on antagonists/agonists on PDGF-C.

25 The transgenic DNA used in any embodiment of the method of the invention comprises a sequence of polynucleotides coding for PDGF-C or an analog thereof, or a functional activated fragment of PDGF-C or an analog thereof. The polynucleotides coding for PDGF-C or an  
30 analog thereof, or a functional activated fragment of PDGF-C or an analog thereof may have been modified in a number of ways known in the art but still retain all or part of the biological activity of PDGF-C.

The "modified polynucleotides coding for PDGF-C" of this invention refers to a sequence of polynucleotides coding for PDGF-C or an analog thereof, or a functional activated fragment of PDGF-C or an analog thereof that has  
5 been modified by an *in vitro* or recombinant DNA technique. Modifications including deletions, substitutions, and insertions of nucleotides into the polynucleotide sequence are specifically included.

Preferably where amino acid substitution is used, the  
10 substitution is conservative, i.e. an amino acid is replaced by one of similar size and with similar charge properties.

Conservative substitution denotes the replacement of an amino acid residue by another, biologically similar  
15 residue. Examples of conservative substitutions include the substitution of one hydrophobic residue such as isoleucine, valine, leucine, alanine, cysteine, glycine, phenylalanine, proline, tryptophan, tyrosine, norleucine or methionine for another, or the substitution of one polar  
20 residue for another, such as the substitution of arginine for lysine, glutamic acid for aspartic acid, or glutamine for asparagine, and the like. Neutral hydrophilic amino acids which can be substituted for one another include asparagine, glutamine, serine and threonine. Conservative  
25 substitution also includes the use of a substituted amino acid in place of an unsubstituted parent amino acid.

As such, it should be understood that in the context of the present invention, a conservative substitution is  
30 recognized in the art as a substitution of one amino acid for another amino acid that has similar properties. Exemplary conservative substitutions are set out in the following Table A from WO 97/09433.

Table A  
Conservative Substitutions I

	SIDE CHAIN	
5	CHARACTERISTIC	AMINO ACID
	Aliphatic	
	Non-polar	G A P I L V
	Polar - uncharged	C S T M
10		N Q
	Polar - charged	D E K R
	Aromatic	H F W Y
	Other	N Q D E
15		

Alternatively, conservative amino acids can be grouped as described in Lehninger, [Biochemistry, Second Edition; Worth Publishers, Inc. NY:NY (1975), pp.71-77] as set out in the following Table B.

Table B

## Conservative Substitutions II

5	SIDE CHAIN CHARACTERISTIC	AMINO ACID
	Non-polar (hydrophobic)	
	A. Aliphatic:	A L I V P
	B. Aromatic:	F W
10	C. Sulfur-containing:	M
	D. Borderline:	G
	Uncharged-polar	
	A. Hydroxyl:	S T Y
	B. Amides:	N Q
15	C. Sulfhydryl:	C
	D. Borderline:	G
	Positively Charged (Basic):	K R H
	Negatively Charged (Acidic):	D E

20 Exemplary conservative substitutions are set out in  
the following Table C.



Table C  
Conservative Substitutions III

	<u>Original</u> <u>Residue</u>	<u>Exemplary</u> <u>Substitution</u>
5	Ala (A)	Val, Leu, Ile
	Arg (R)	Lys, Gln, Asn
	Asn (N)	Gln, His, Lys, Arg
	Asp (D)	Glu
	Cys (C)	Ser
10	Gln (Q)	Asn
	Glu (E)	Asp
	His (H)	Asn, Gln, Lys, Arg
	Ile (I)	Leu, Val, Met, Ala, Phe,
	Leu (L)	Ile, Val, Met, Ala, Phe
15	Lys (K)	Arg, Gln, Asn
	Met (M)	Leu, Phe, Ile
	Phe (F)	Leu, Val, Ile, Ala
	Pro (P)	Gly
	Ser (S)	Thr
20	Thr (T)	Ser
	Trp (W)	Tyr, Phe
	Tyr (Y)	Trp, Phe, Thr, Ser
	Val (V)	Ile, Leu, Met, Phe, Ala

25        In yet another aspect, the invention provides a method for screening compounds for the ability to inhibit the effects of PDGF-C, such as hypertrophy or fibrosis of the

heart, liver, kidney, pancreas, ovaries, and other tissues. The method comprises introducing the compound into a transgenic animal comprising cells which overexpress PDGF-C and monitoring, by any suitable means, an inhibition in the biological activity of PDGF-C in the animal. Monitoring includes but is not limited to comparing the wildtype and transgenic animals.

10 Brief Description of the Drawings

The invention will be described in further detail hereinafter with reference to the accompanying drawings in which:

15 Figure 1 shows body weights of normal (wild type, wt) and transgenic (tg) mice at one and three months of age.

Figure 2 shows heart weights of normal and transgenic mice at one and three months of age.

Figure 3A shows a tissue section from a normal (wild type, wt) mouse heart.

20 Figure 3B shows a tissue section from a transgenic (tg) mouse heart.

Figure 4 shows the results of SDS-PAGE and immunoblotting of tissue extracts from transgenic (tg) and normal (wild type, wt) mouse hearts.

25 Figure 5 shows a normal (wild type, wt, to the left) and a transgenic (tg, to the right) mouse heart from 6 month old male mice.

30 Figure 6A shows microvessel organization in a stained heart left ventricle section from a normal (wild type, wt) six month old mouse.

Figure 6B shows microvessel disorganization in a stained heart left ventricle section from a transgenic (tg) six month old mouse.

Detailed Description of Illustrative Embodiments

The descriptions and examples below are exemplary of the embodiments and scope of this invention. The invention is not limited to the scope of this description.

5       Persons skilled in the art will appreciate that the following examples and embodiments may be modified using techniques known in the art. For example, variations in the nucleic acid sequences described or claimed can be produced by known methods without altering the effects or  
10       advantages the inventors have shown. Such variations are therefore included within the scope of this description and invention. In addition, detailed protocols for many of the techniques known in the art are described by Ausubel, F.M. et al. Eds. *Current Protocols in Molecular Biology*,  
15       Greene Publishing Associates and Wiley-Interscience, John Wiley & Sons, Boston, MA (1989), and Supplements through Jan. 1997; in Sambrook, J., et al., *Molecular Cloning. A Laboratory Manual*, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989); in B. Hogan et al.  
20       Eds. *Manipulating the Mouse Embryo. A Laboratory Manual*, Cold Spring Harbor Laboratory Press, (1994); and in Gordon and Ruddle, *Science*, 1981 214 1244-1246. These documents are specifically incorporated herein by reference, and may be relied on to evidence enablement of one skilled in the  
25       art to practice the embodiments of the invention.

Transgenic DNA refers to DNA that is introduced into a cell so that the DNA is incorporated into the genome of the cell. The cell may be capable of giving rise to a transgenic animal which contains the transgenic DNA.  
30       Generally, the transgenic DNA for administration into a particular cell can be constructed using a transgenic vector. A preferred DNA is a polynucleotide that encodes for full-length PDGF-C or an analog thereof, and a more

preferred DNA is that which encodes for the activated truncated PDGF-C or an analog thereof. The truncated portion of PDGF-C comprises at least a portion of the PDGF/VEGF homology domain (P/VHD) of PDGF-C. The minimal  
5 sequence is residues 230-245 of SEQ ID NO:1. However, the domain can extend towards the N terminus up to residue 164 of SEQ ID NO:1. Herein the P/VHD of PDGF-C is defined as truncated PDGF-C. The truncated PDGF-C is an activated form of PDGF-C.

10 A recombinant gene or sequence simply means the gene or sequence has been manipulated in any one of a number of recombinant DNA techniques known in the art.

As used herein the term "modified polynucleotide sequence coding for PDGF-C" refers to a PDGF-C  
15 polynucleotide sequence from an animal which has been modified by one or more of: point mutations, site-directed mutagenesis, deletions, and insertions. Alternatively, a modified polynucleotide sequence coding for PDGF-C is a sequence which is linked to a second sequence, such as a  
20 marker sequence, epitope tag or promoter/enhancer, which is not associated with wild type PDGF-C DNA. PDGF-C encoded by a modified polynucleotide sequence retains some or all of the activities of wild type PDGF-C, for example, the PDGFR-alpha binding motif.

25 The FLAG™ peptide can be used as an epitope tag in many cell types. The sequence, use and detection of the FLAG™ tag is described in Chubet, RG, et. al., Vectors for expression and secretion of FLAG epitope-tagged proteins in mammalian cells, *Biotechniques* 1996 Jan;20(1):136-41.  
30 Constructing a mammalian expression vector using, for example GAL4 and a Flag peptide is described in Witzgall, R., et. al., A mammalian expression vector for the

expression of GAL fusion proteins with an epitope tag and histidine tail, *Anal Biochem* 1994 Dec;223(2):291-8.

As noted above, there are many known processes for generating transgenic animals. These processes are essentially the same regardless of the species involved. While the Examples that follow describe transgenic mice, the same techniques may be used to produce non-mouse transgenic animals, and their creation and use is encompassed within the scope of this invention.

One process begins with transgenic DNA operably linked to a promoter. The transgenic DNA-promoter complex is introduced into the pronuclei of a fertilized egg of a non-human animal. The egg is then implanted into a pseudopregnant non-human animal and allowed to develop into a transgenic animal.

Fertilized eggs from a variety of animals used in the above described method can be produced using techniques well known to those of ordinary skill in the art. For example, the use of bovine oocytes to support embryos of a number of species is described in Dominko, T., et. Al., *Bovine oocyte cytoplasm supports development of embryos produced by nuclear transfer of somatic cell nuclei from various mammalian species*, *Biol Reprod*, 1999 Jun;60(6):1496-502.

Alternatively, fertilized eggs from a variety of animals can be obtained from a number of sources. These various species include mice, cows, rabbits, and sheep, as well as other animals (Mullins et al, *J. Clin. Invest.*, 1996 98 S37-S40). Accordingly, the invention is as applicable to animals other than the specifically exemplified mice. While many of the references regarding the state of the art relate to mammalian species, the

invention is applicable and enabled to animals other than mammals.

A second method for producing transgenic animals involves the modification of embryonic stem (ES) cells. This second method comprises introducing transfected cells into embryos at a stage at which they are capable of integrating into the embryo, for example, at the blastocyte stage. The embryo with transfected cells is then replanted into a surrogate mother, resulting in chimeric offspring possessing the transgenic DNA.

Embryonic stem cells are available from a number of sources. These include mice, rats, cows, pigs, sheep, and other animals (Joyner A.L., (1993), *Gene Targeting*). Alternatively, the production of ES cells from a variety of animals is well known to those of ordinary skill in the art. A practical approach, edited by Wood, R. and Hames, B.D., *The Practical Approach Series*, vol. 126, Oxford IRL Press (specifically incorporated herein by reference) describes methods of producing ES cells. Also, B. Hogan et al., Eds., *Manipulating the Mouse Embryo. A laboratory Manual*, Cold Spring Harbor Laboratory Press, (1994) describes manipulating the mouse embryo.

Methods used in successful research with rat ES cells are described in Iannaccone, PM, et. al., Pluripotent embryonic stem cells from the rat are capable of producing chimeras, *Dev Biol*, 1994 May;163(1):288-92. Work with rabbit ES cells is described in Schoonjans, L., et. al., Pluripotential rabbit embryonic stem (ES) cells are capable of forming overt coat color chimeras following injection into blastocysts, *Mol Reprod Dev*, 1996 Dec;45(4):439-43. In addition, Couly and Le Douarin, *Development*, 1990 108 543-555, describes methods for isolating and manipulating

chicken and quail embryos. Kimmel and Warga, *Nature*, 1987 327 234-237, describe isolation and manipulation of zebrafish embryos. Ware et al., "Development of Embryonic Stem Cell Lines from Farm Animals," *Society for the Study of Reproduction*, 1988 38 241 also discusses an embryonic stem cell culture condition amenable for many species like mouse, cattle, pig, and sheep.

Specific references for pig embryonic stem cells include Notarianni E., et. al., Incorporation of cultured embryonic cells into transgenic and chimeric, porcine fetuses, *Int J Dev Biol*, 1997 Jun;41(3):537-40 and Gutierrez-Adan, et. al., Isolation of Pluripotent Stem Cells from Cultured Porcine Primordial Germ Cells, *Biol Reprod*, Nov;57(5):1089-95 (1997). Specific references for cows are Cibelli, et. al., Transgenic Bovine Chimeric Offspring Produced from Somatic Cell-Derived Stem-Like Cells, *Nat. Biotechnol*, Jul;16(7):642-6 (1998) and Kubota C., et. al., Six cloned calves produced from adult fibroblast cells after long-term culture, *Proc Natl Acad Sci USA*, 2000 Feb 1;97(3):990-5. Preparing primate embryonic stem cells can be facilitated by referring to Thompson, et. al., Isolation of a Primate Embryonic Stem Cell Line, *Proc. Natl Acad Sci USA*, Aug 15;92(17):7844-8 (1995).

Various methods are known in the art for introducing DNA into animal cells, for example, ES cells. Transgenic DNA can be microinjected into the appropriate cells. Also, viral vectors can be used to introduce the DNA into appropriate cells and the genome of those cells (See, for example, Tsukui et al., *Nature Biotechnology*, 1996 14 982-985). Or, cells can be manipulated *in vitro* through transfection and electroporation methods (See, for example,

Ausubel, F.M. et al. Eds. *Current Protocols in Molecular Biology*, Greene Publishing Associates and Wiley-Interscience, John Wiley & Sons, Boston, MA (1989); and B. Hogan et al., Eds., *Manipulating the Mouse Embryo. A Laboratory Manual*, Cold Spring Harbor Laboratory Press, (1994)).

Generally, this type of transgenic DNA incorporates into a cell genome through random integration, although homologous recombination is possible. The design of transgenic DNA vectors involves linking the transgenic DNA to an appropriate promoter sequence. Examples of promoters that can be used include, but are not limited to, alpha-myosin heavy chain promoter which gives cardiac myocyte-specific expression, keratin K14 promoter which gives basal keratinocyte-specific expression, and insulin promoter which gives pancreatic beta cell-specific expression.

#### Examples

Mice overexpressing PDGF-C were created to analyze the role of PDGF-C in development and maintenance *in vivo* under diseased states. The transgenic mice were constructed to overexpress a c-myc epitope-tagged PDGF-C using the alpha-myosin heavy chain (alpha-MHC) promoter (Subramaniam, A. et al., J. Biol. Chem. 1991 266 24613-24620).

#### Example 1: Insertion of the Transgenic Vector Into an Animal Cell.

The sequence encoding the human c-myc epitope (Evan, G.I. et al., Mol. Cell Biol., 1985 5 3610-3616) was introduced at the 3' end of the coding region of mouse PDGF-C cDNA by PCR mutagenesis with DeepVent polymerase



(Biolabs). The primers used for PCR were 5'-CGGAATTCTCAG  
CCAAATGCTCCTCCTC (forward) (SEQ ID NO:3) and 5'-CGGATTTCTTAC  
AAGTCTTCTTCAGAAATAAGCTTTTGTTCCTCCTGCGTTTCCTCT (reverse)  
(SEQ ID NO:4). This generated a 1100 bp fragment which was  
5 subcloned into the pBluescript-based transgenic vector  
between the 5.5 kb mouse alpha-myosin heavy chain promoter  
and the 250 bp polyadenylation sequence from SV-40  
(Subramaniam, A., et al., J. Biol. Chem. 1991 266 24613-  
24620). The linearized and purified transgene fragment was  
10 microinjected into the male pronuclei of fertilized mouse  
oocytes (Mouse Camp, Karolinska Institutet, Stockholm).

Example 2: Production of Transgenic Animals.

The injected fertilized oocytes were implanted into  
15 pseudopregnant foster mothers. Tail DNA was collected from  
the resulting heterozygous pups. Generally, 0.5 cm of tail  
tissue was surgically removed and used to prepare DNA  
samples. The tail DNA was screened for the presence of the  
transgene by PCR using the mouse PDGF-C specific primer,  
20 5'-GTGTCCATACGGGAAGAG (forward) (SEQ ID NO:5) and the human  
c-myc specific primer 5'-GTCTTCTTCAGAAATAAGC (reverse) (SEQ  
ID NO:6). A 294 bp fragment was amplified in the  
transgenic founders. Male founders were backcrossed to  
C57Bl/6 wildtype females.

25

Example 3: Comparative body and heart weights at one and  
three months of age.

Body and heart weights were measured at one and three  
30 months of age for both normal (wild type, wt) and  
transgenic (tg) mice. Live mice were weighed on a normal  
laboratory scale and heart weights were obtained after  
dissection of the hearts.

The weights were sorted by type (normal or transgenic), sex (male or female) and age (one or three months). The data for body weight are summarized in Figure 1. A general reduction in body weight is seen in the transgenic animals. The data for heart weight are summarized in Figure 2. Heart weights increase in the transgenic animals, particularly in the male mice.

Example 4: Comparative heart studies at six to eight weeks of age.

At six to eight weeks of age, transgenic and normal animals were killed and their hearts removed, washed in ice cold PBS and prepared for routine histology and biochemical analysis. Hearts were fixed in 4% paraformaldehyde in PBS overnight at 4°C, dehydrated, embedded in paraffin wax and sectioned. The tissue sections were stained in hematoxylin/eosin using standard techniques.

Analysis of tissue sections from hearts of adult normal (wild type, wt) (Figure 3A) and transgenic (tg) animals (Figure 3B) showed that overexpression of PDGF-C induced a strong proliferation of myocardial interstitial cells, e.g. cardiac fibroblasts. Cardiac fibroblasts express PDGFR-alpha and PDGF-AA dimer has previously been shown to be a potent mitogen for this cell type (Simm, A. et al., *Basic Res. Cardiol.* 1998 93 Suppl 3: 40-43). The expansion of the interstitium in the transgenic hearts caused a drastic disorganization of the cardiac myofibers indicating that the functional properties of the transgenic hearts were severely compromised.

For biochemical analysis, hearts were minced and then sonicated in ice cold 10 mM Tris-HCl buffer pH 7.6 containing 0.1M NaCl, 1mM EDTA, 1mg/ml aprotinin and 4 mM phenylmethylsulfonyl fluoride. Supernatants were collected

after centrifugation at 12,000 g for 20 min at 4°C, and aliquots were subsequently subjected to SDS-PAGE under reducing conditions. For immunoblotting analysis anti-c-myc mouse ascites (9E10) and a rabbit anti-mouse PDGF-C peptide antiserum were used. The rabbit anti-mouse PDGF-C peptide antiserum was generated using the peptide CVKKSRVVNLNLL KEEVKLYSC (SEQ ID NO:7) (residues 230-250 of mouse PDGF-C (SEQ ID NO:2)) as described in International Application No. PCT/US99/22668. Bound antibodies were visualized using the ECL technique.

Figure 4 shows the results of the analysis of SDS-PAGE and immunoblotting of tissue extracts from normal and transgenic mouse hearts. Transgenic c-myc epitope-tagged PDGF-C was visualized using a c-myc-specific monoclonal antibody (c-myc MAb). Analysis under reducing conditions showed abundant expression of full-length 52 kDa c-myc-tagged PDGF-C in transgenic hearts and the presence of several processed intermediates with a 34 kDa species being most prominent. Similar analysis using the anti-peptide antiserum to PDGF-C confirmed these results. The analysis indicates that proteases able to convert overexpressed latent full-length PDGF-C *in vivo* are expressed in the myocardium, and that activated PDGF-C is able to promote proliferation of PDGFR-alpha expressing cardiac fibroblasts.

The heart phenotype induced in the PDGF-C transgenic animals, with an expansion of the cardiac interstitium, is reminiscent of myocardial hypertrophy and fibrosis. Cardiac fibroblasts synthesize the extracellular matrix and play a pivotal role in adaptation and remodelling of the interstitium, typically seen following myocardial infarctions and increased hemodynamic load. Given that PDGF-C is normally expressed in heart, this indicates that

PDGF-C has a role in physiological and pathophysiological remodelling of the cardiac interstitium.

As cardiac fibrosis is a common consequence following heart infarction, an animal model can be used both to understand the mechanisms underlying this phenomena and as a research tool for developing drugs that may interfere with development of hypertrophy and cardiac fibrosis.

Example 5: Comparative heart gross examinations at six months of age.

Hearts were obtained from normal and transgenic male mice at age six months. Figure 5 shows a normal (wild type, wt) heart on the left and a transgenic (tg) heart on the right. There is an observable general increase in size of the transgenic mouse heart, with a prominent enlargement of the atrial region.

Example 6: Comparative heart histological evaluations at six months of age.

Hearts were collected from six month old mice. The hearts were prepared in paraffin-embedded fixed tissue blocks and regular (5µm) sections were taken. The sections were rehydrated and treated with trypsin (1x TRYPSIN-EDTA, LIFE TECHNOLOGIES) for 20 minutes at 37°C. The sections were then stained for PECAM-1 expression, and endothelial cell marker, using TSA™-Indirect as suggested by the supplier (NEN™ Life Science Products). The primary antibody, rat α PECAM-1 antibody (PharMingen), was incubated at a dilution of 1:500 and the secondary antibody, biotinylated rat α IgG (mouse adsorbed, Vector Laboratories), was incubated at a dilution of 1:300. The chromogenic substrate used was 3,3'-DAB (Sigma).

Photographs of the left ventricles were taken and are reproduced in Figures 6A and 6B. It can be observed that microvessel organization in the wild-type (wt, Figure 6A) mouse heart shows an even distribution of capillaries. In contrast, the disorganized microvessels evident in the transgenic mouse heart (Figure 6B) show a loss of capillaries and the formation of unevenly distributed vascular sac-like structures. The formation of these sac-like structures is likely a sign of proliferation of the vasculature. Sac-like structure formation may be an indirect sign of insufficient vascular supply to the heart. This type of microvessel disease is frequently observed in human disease, particularly in heart fibrosis.

Example 7: Comparative heart histological evaluations.

Further analysis was conducted by staining wild type and transgenic hearts with Mason trichromate to show deposition of extracellular matrix. Regular sections from paraformaldehyde-fixed and paraffin-embedded tissue blocks were deparaffinized and hydrated with 2 x xylene for 5 minutes, 2 x abs for 3 minutes, 2 x 95% for 3 minutes, 70% for 5 minutes, followed by dH<sub>2</sub>O. Bouin's solution (a mixture of 75 ml saturated aqueous picric acid, 25 ml of concentrated (37-40%) formalin, and 5 ml of glacial acetic acid) was preheated to 58°C and the sections were incubated in the solution for 15 minutes. The sections were washed in running water approximately 4-10 minutes, until the yellow color disappeared, then rinsed twice with dH<sub>2</sub>O.

A working solution of Modified Weighert's iron hematoxylin was prepared using equal parts of solution A and solution B and stored protected from light. Solution A consists of 2 grams of hematoxylin crystals (C.I. 75290) and 100 ml of 90% alcohol. Solution B consists of 4 ml of

62% aqueous ferric chloride ( $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$ ), 95 ml of  $\text{dH}_2\text{O}$ , and 1 ml of concentrated  $\text{HCl}$ .

After the proceeding steps, the sections were placed in the Modified Weigert's solution for 2 minutes, then washed in running water and rinsed twice with  $\text{dH}_2\text{O}$ . Next, Biebrich scarlet-acid fuchsin (90 ml of 1% aqueous Biebrich scarlet (C.I. 26905) with 10 ml of 1% aqueous acid fuchsin (C.I. 42685) and 1 ml glacial acetic acid, stored protected from light) was applied for a few seconds, followed by three rinses with  $\text{dH}_2\text{O}$ . The sections were placed in phosphomolybdic-phosphotungstic acid solution (2.5 g of phosphomolybdic acid, 2.5 g of phosphotungstic acid, and 100 ml  $\text{dH}_2\text{O}$ ) for 5 minutes, followed by 30 minutes in water blue solution (3 g of water blue, 2 ml of glacial acetic acid, and 100 ml  $\text{dH}_2\text{O}$ , stored protected from light). The sections were carefully rinsed with  $\text{dH}_2\text{O}$ . Acetic acid solution (1 ml glacial acetic acid with 100 ml  $\text{dH}_2\text{O}$ ) was applied for 3 minutes, followed by two rinses with  $\text{dH}_2\text{O}$ .

The sections were then dehydrated in alcohol series to histoclear and mounted with entellan. The sections were allowed to dry in a hood. The staining yielded black nuclei, blue collagen, and red cytoplasm, keratin, muscle fibers, and fibrin.

This procedure revealed staining in hearts from one month old transgenic animals similar to staining in the hearts of wild type animals. However, the phenotype gets progressively worse with age. At three months of age, hearts from transgenic animals show significant deposition of the extracellular matrix. At six months of age, hearts from transgenic animals show extensive deposition of the extracellular matrix.

This example demonstrates PDGF-C overexpression in transgenic animals causes proliferation of cardiac

fibroblasts and a resultant expansion of the interstitium. The overexpression of PDGF-C may also induce the fibroblasts to secrete more extracellular matrix, essentially collagens. This example provides further  
5 evidence that the overexpression of PDGF-C induces fibrosis.

Example 8: Method for PDGF-C antagonist compound identification in an animal system.

10 A compound is identified as a PDGF-C antagonist by introducing the compound into a transgenic, non-human animal which overexpresses PDGF-C or an analog thereof, or a functional fragment of PDGF-C or an analog thereof. Following introduction of the compound, biological activity  
15 of PDGF-C in the animal is monitored. An inhibition of PDGF-C biological activity indicates the compound may be useful as a PDGF-C antagonist. Comparing biological activity of PDGF-C could involve comparing the transgenic animal with a normal (wild type) animal of the same  
20 species.

Example 9: Method for PDGF-C antagonist compound identification in a cellular system.

25 A compound is identified as a PDGF-C antagonist by introducing the compound into a cell or cells isolated from a transgenic, non-human animal which overexpresses PDGF-C or an analog thereof, or a functional fragment of PDGF-C or an analog thereof. Following introduction of the compound, biological activity of PDGF-C in the cell or cells is  
30 monitored. An inhibition of PDGF-C biological activity indicates the compound may be useful as a PDGF-C antagonist.

Example 10: Method for screening compounds targeted to hypertrophy.

A compound may be evaluated for its ability to inhibit hypertrophy by administering a pharmaceutically active amount of the compound to a transgenic, non-human animal which overexpresses PDGF-C or an analog thereof, or a functional fragment of PDGF-C or an analog thereof. Following introduction of the compound, the animal's cardiac development is monitored. Normal cardiac development in the animal may indicate the compound is useful in the inhibition of hypertrophy.

Example 11: Method for screening compounds targeted to fibrosis.

A compound may be evaluated for its ability to inhibit fibrosis by administering a pharmaceutically active amount of the compound to a transgenic, non-human animal which overexpresses PDGF-C or an analog thereof, or a functional fragment of PDGF-C or an analog thereof. Following introduction of the compound, the animal's cardiac development is monitored. Normal cardiac development in the animal may indicate the compound is useful in the inhibition of fibrosis.

The foregoing description and examples have been set forth merely to illustrate the invention and are not intended to be limiting. Since modifications of the disclosed embodiments incorporating the spirit and substance of the invention may occur to persons skilled in the art, the invention should be construed to include everything within the scope of the appended claims and equivalents thereof. All references cited herein are expressly incorporated by reference.



What is claimed is:

1. A method for producing a transgenic, non-human animal overexpressing PDGF-C or an analog thereof, or a functional fragment of PDGF-C or an analog thereof, the method comprising the steps of:

a) introducing a transgenic DNA into a cell of a non-human animal, said transgenic DNA comprising a polynucleotide sequence encoding for PDGF-C or an analog thereof, or a functional fragment of PDGF-C or an analog thereof;

b) allowing said transgenic DNA to integrate into said cell;

c) introducing said cell from step b) into a non-human animal; and

d) allowing said cell from step c) to develop into a transgenic, non-human animal.

2. A method for producing a transgenic, non-human animal overexpressing PDGF-C or an analog thereof, or a functional fragment of PDGF-C or an analog thereof, the method comprising the steps of:

a) introducing a transgenic DNA into the pronuclei of a fertilized oocyte of a non-human animal, said transgenic DNA comprising a polynucleotide sequence encoding for PDGF-C or an analog thereof, or a functional fragment of PDGF-C or an analog thereof;

b) implanting said fertilized oocyte into a pseudopregnant non-human animal; and

c) allowing said fertilized oocyte to develop into a transgenic, non-human animal.

3. A method for producing a transgenic, non-human animal overexpressing PDGF-C or an analog thereof, or a functional fragment of PDGF-C or an analog thereof, the method comprising the steps of:

- a) introducing a transgenic DNA into an embryonic stem cell of a non-human animal, said transgenic DNA comprising a polynucleotide sequence encoding for PDGF-C or an analog thereof, or a functional fragment of PDGF-C or an analog thereof;
- b) selecting an embryonic stem cell when said transgenic DNA has integrated into the genomic DNA of said embryonic stem cell;
- c) introducing the selected cell from step b) into an embryo of a developing non-human animal; and
- d) allowing said embryo to develop into a transgenic, non-human animal.

4. The method of any of claims 1-3, wherein said transgenic DNA is operably linked to a promoter.

5. The method of claim 4, wherein said promoter is selected from group consisting of: alpha-myosin heavy chain promoter, keratin K14 promoter, and insulin promoter.

6. The method of any of claims 1-3, wherein said transgenic DNA is operably linked to an epitope-tag.

7. The method of claim 6, wherein said epitope tag is c-myc.

8. The method of any of claims 1-3, wherein said transgenic DNA is operably linked to a marker sequence.

9. A transgenic, non-human animal produced by the method of any of any of claims 1-3.

10. An animal according to claim 9, wherein said animal is a rodent.

11. An animal according to claim 10, wherein said animal is a mouse.

12. A transgenic, non-human animal that is a descendant from an animal according to claim 9.

13. A transgenic, non-human animal that is a descendant from an animal according to claim 10.

14. A transgenic, non-human animal that is a descendant from an animal according to claim 11.

15. A cell isolated from an animal according to claim 9.

16. A cell isolated from an animal according to claim 10.

17. A cell isolated from an animal according to claim 11.

18. A fertilized oocyte containing transgenic DNA that encodes for PDGF-C or an analog thereof, or a functional fragment of PDGF-C or analog thereof.

19. An embryonic stem cell containing transgenic DNA that encodes for PDGF-C or an analog thereof, or a functional fragment or analog thereof.

20. A method for identifying a compound as a PDGF-C antagonist, said method comprising the steps of:

introducing said compound into a transgenic, non-human animal overexpressing PDGF-C or an analog thereof, or a functional fragment of PDGF-C or an analog thereof;

monitoring the biological activity of PDGF-C in said animal; and

identifying said compound as a PDGF-C antagonist where PDGF-C biological activity is inhibited.

21. The method of claim 20 wherein said monitoring step comprises comparing said transgenic, non-human animal with a wild-type non-human animal of the same species.

22. A method for identifying a compound as a PDGF-C antagonist, said method comprising the steps of:  
introducing said compound into a cell isolated from a transgenic, non-human animal overexpressing PDGF-C or an analog thereof, or a functional fragment of PDGF-C or an analog thereof;  
assaying the effect of said compound on said cell; and  
identifying said compound as a PDGF-C antagonist where the PDGF-C biological activity of said cell is altered.

23. A method of screening a compound for inhibition of hypertrophy, comprising the steps of:  
administering a pharmaceutically active amount of said compound to a transgenic, non-human animal overexpressing PDGF-C or an analog thereof, or a functional fragment of PDGF-C or an analog thereof; and  
monitoring the cardiac development of said animal;  
determining said compound inhibits hypertrophy where said cardiac development is normal.

24. A method of screening a compound for inhibition of fibrosis, comprising the steps of:  
administering a pharmaceutically active amount of said compound to a transgenic, non-human animal overexpressing PDGF-C or an analog thereof, or a functional fragment of PDGF-C or an analog thereof; and  
monitoring the cardiac development of said animal;  
determining said compound inhibits fibrosis where said cardiac development is normal.

1/3

### Body Weight




Figure 1

### Heart Weight




Figure 2

2/3



Figure 3A



Figure 3B





Figure 4

3/3



10 mm  
Figure 5




Figure 6A



20 μm  
Figure 6B

SEQUENCE LISTING

<110> ERIKSSON, Ulf

LI, Xuri

PONTEN, Annica

AASE, Karin

LI, Hong

<120> NON-HUMAN TRANSGENIC ANIMALS EXPRESSING PLATELET-DERIVED  
GROWTH FACTOR C (PDGF-C) AND USES THEREOF

<130> 1064/48487

<150> US 60/192,507

<151> 2000-03-28

<160> 7

<170> PatentIn version 3.0

<210> 1

<211> 345

<212> PRT

<213> Homo sapiens

<400> 1

Met Ser Leu Phe Gly Leu Leu Leu Val Thr Ser Ala Leu Ala Gly Gln  
1            5            10            15



Arg Arg Gly Thr Gln Ala Glu Ser Asn Leu Ser Ser Lys Phe Gln Phe  
20 25 30

Ser Ser Asn Lys Glu Gln Asn Gly Val Gln Asp Pro Gln His Glu Arg  
35 40 45

Ile Ile Thr Val Ser Thr Asn Gly Ser Ile His Ser Pro Arg Phe Pro  
50 55 60

His Thr Tyr Pro Arg Asn Thr Val Leu Val Trp Arg Leu Val Ala Val  
65 70 75 80

Glu Glu Asn Val Trp Ile Gln Leu Thr Phe Asp Glu Arg Phe Gly Leu  
85 90 95

Glu Asp Pro Glu Asp Asp Ile Cys Lys Tyr Asp Phe Val Glu Val Glu  
100 105 110

Glu Pro Ser Asp Gly Thr Ile Leu Gly Arg Trp Cys Gly Ser Gly Thr  
115 120 125

Val Pro Gly Lys Gln Ile Ser Lys Gly Asn Gln Ile Arg Ile Arg Phe  
130 135 140

Val Ser Asp Glu Tyr Phe Pro Ser Glu Pro Gly Phe Cys Ile His Tyr  
145 150 155 160

Asn Ile Val Met Pro Gln Phe Thr Glu Ala Val Ser Pro Ser Val Leu  
165 170 175

Pro Pro Ser Ala Leu Pro Leu Asp Leu Leu Asn Asn Ala Ile Thr Ala  
180 185 190

Phe Ser Thr Leu Glu Asp Leu Ile Arg Tyr Leu Glu Pro Glu Arg Trp  
195 200 205

Gln Leu Asp Leu Glu Asp Leu Tyr Arg Pro Thr Trp Gln Leu Leu Gly  
210 215 220

Lys Ala Phe Val Phe Gly Arg Lys Ser Arg Val Val Asp Leu Asn Leu  
225 230 235 240

Leu Thr Glu Glu Val Arg Leu Tyr Ser Cys Thr Pro Arg Asn Phe Ser  
245 250 255

Val Ser Ile Arg Glu Glu Leu Lys Arg Thr Asp Thr Ile Phe Trp Pro  
260 265 270

Gly Cys Leu Leu Val Lys Arg Cys Gly Gly Asn Cys Ala Cys Cys Leu  
 275 280 285

His Asn Cys Asn Glu Cys Gln Cys Val Pro Ser Lys Val Thr Lys Lys  
 290 295 300

Tyr His Glu Val Leu Gln Leu Arg Pro Lys Thr Gly Val Arg Gly Leu  
 305 310 315 320

His Lys Ser Leu Thr Asp Val Ala Leu Glu His His Glu Glu Cys Asp  
 325 330 335

Cys Val Cys Arg Gly Ser Thr Gly Gly  
 340 345

<210> 2

<211> 345

<212> PRT

<213> Murinae gen. sp.

<400> 2

Met Leu Leu Leu Gly Leu Leu Leu Thr Ser Ala Leu Ala Gly Gln  
 1 5 10 15

Arg Thr Gly Thr Arg Ala Glu Ser Asn Leu Ser Ser Lys Leu Gln Leu  
 20 25 30

Ser Ser Asp Lys Glu Gln Asn Gly Val Gln Asp Pro Arg His Glu Arg  
 35 40 45

Val Val Thr Ile Ser Gly Asn Gly Ser Ile His Ser Pro Lys Phe Pro  
 50 55 60

His Thr Tyr Pro Arg Asn Met Val Leu Val Trp Arg Leu Val Ala Val  
 65 70 75 80

Asp Glu Asn Val Arg Ile Gln Leu Thr Phe Asp Glu Arg Phe Gly Leu  
 85 90 95

Glu Asp Pro Glu Asp Asp Ile Cys Lys Tyr Asp Phe Val Glu Val Glu  
 100 105 110

Glu Pro Ser Asp Gly Ser Val Leu Gly Arg Trp Cys Gly Ser Gly Thr  
115 120 125

Val Pro Gly Lys Gln Thr Ser Lys Gly Asn His Ile Arg Ile Arg Phe  
130 135 140

Val Ser Asp Glu Tyr Phe Pro Ser Glu Pro Gly Phe Cys Ile His Tyr  
145 150 155 160

Ser Ile Ile Met Pro Gln Val Thr Glu Thr Thr Ser Pro Ser Val Leu  
165 170 175

Pro Pro Ser Ser Leu Ser Leu Asp Leu Leu Asn Asn Ala Val Thr Ala  
180 185 190

Phe Ser Thr Leu Glu Glu Leu Ile Arg Tyr Leu Glu Pro Asp Arg Trp  
195 200 205

Gln Val Asp-Leu Asp Ser Leu Tyr Lys Pro Thr Trp Gln Leu Leu Gly  
210 215 220

Lys Ala Phe Leu Tyr Gly Lys Lys Ser Lys Val Val Asn Leu Asn Leu  
225 230 235 240

Leu Lys Glu Glu Val Lys Leu Tyr Ser Cys Thr Pro Arg Asn Phe Ser  
245 250 255

Val Ser Ile Arg Glu Glu Leu Lys Arg Thr Asp Thr Ile Phe Trp Pro  
260 265 270

Gly Cys Leu Leu Val Lys Arg Cys Gly Gly Asn Cys Ala Cys Cys Leu  
275 280 285

His Asn Cys Asn Glu Cys Gln Cys Val Pro Arg Lys Val Thr Lys Lys  
290 295 300

Tyr His Glu Val Leu Gln Leu Arg Pro Lys Thr Gly Val Lys Gly Leu  
305 310 315 320

His Lys Ser Leu Thr Asp Val Ala Leu Glu His His Glu Glu Cys Asp  
325 330 335

Cys Val Cys Arg Gly Asn Ala Gly Gly  
340 345

<210> 3

<211> 28

<212> DNA

<213> Murinae gen. sp.

<400> 3

cggaattctc agccaaatgc tcctcctc

28

<210> 4

<211> 59

<212> DNA

<213> Murinae gen. sp.

<220>

<221> unsure

<222> (1)..(59)

<223> Also contains sequence encoding the human c-myc epitope

<400> 4

cggatttctt acaagtcttc ttcagaaata agcttttgtt cccctcctgc gtttcctct 59

<210> 5

<211> 18

<212> DNA

<213> Murinae gen. sp.

<400> 5

gtgtccatac gggaagag

18

&lt;210&gt; 6

&lt;211&gt; 19

&lt;212&gt; DNA

&lt;213&gt; Murinae gen. sp.

&lt;400&gt; 6

gtcttcttca gaaataagc

19

&lt;210&gt; 7

&lt;211&gt; 22

&lt;212&gt; PRT

&lt;213&gt; Murinae gen. sp.

&lt;220&gt;

&lt;221&gt; UNSURE

&lt;222&gt; (1)..(22)

&lt;223&gt; Residues 230-250 of mouse PDGF-C (SEQ ID NO:2)

&lt;400&gt; 7

Cys Val Lys Lys Ser Arg Val Val Asn Leu Asn Leu Leu Lys Glu Glu  
1            5            10            15

Val Lys Leu Tyr Ser Cys  
20

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US01/09855

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : Please See Extra Sheet.  
US CL : 424/93.1, 93.21; 800/3, 9, 18, 24, 25

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
U.S. : 424/93.1, 93.21; 800/3, 9, 18, 24, 25

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)  
Please See Extra Sheet.

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5,434,132A (ROZENGURT et al) 18 July 1995, see whole document.	1-24
A	DING et al. The Mouse Pdgfc Gene: Dynamic Expression in Embryonic Tissues During Organogenesis. Mech. Dev. 2000. Vol 96. pages 209-213.	1-24
A	LI et al. PDGF-C is a New Protease-Activated Ligand for the PDGF alpha Receptor. Nature Cell Biology. May 2000. Vol. 2. pages 302-309.	1-24

☒ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

* Special categories of cited documents:	* "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier document published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search  
30 MAY 2001

Date of mailing of the international search report

21 JUN 2001

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Authorized officer

BRIAN WHITEMAN

Facsimile No. (703) 305-3230

Telephone No. (703) 308-0196

Form PCT/ISA/210 (second sheet) (July 1998) \*

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US01/09855

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	SHANKAR et al. Thrombin Stimulates PDGFC Production by Human Aortic Endothelial Cells through Activation of a G Protein and the Na <sup>+</sup> /H <sup>+</sup> Antiporter. J. Cell. Biology. November 1989. Vol. 109 (4 part 2). page 152A.	1-24
A	ZWERNER et al. PDGF-C is an EWS/FLI Induced Transforming Growth Factor in Ewing Family Tumors. Oncogene. 2001. Vol. 20. pages 626-633.	1-24
A	BERGSTEN et al. PDGF-D is a Specific, Protease-Activated Ligand for the PDGF beta-Receptor. Nature Cell Biology. May 2001. Vol. 3. pages 512-516.	1-24
A	VILOTTE et al. Modification and Repression of Genes Expressed in the Mammary Gland Using Gene Targeting and Other Technologies. J. Mammary Gland Biol. and Neoplasia. July 1998. Vol. 3. No. 3. pages 351-362.	1-20
A	NOMURA, T. Practical Development of Genetically Engineered Animals as Human Disease Models. Laboratory Animal Science. April 1997. Vol. 47. No. 2. pages 113-117.	1-20
A	MULLINS et al. Perspective Series: Molecular Medicine in Genetically Engineered Animals. J. Clin. Invest. 1996. Vol. 98. No. 11. Supplement. pages S37-S40.	1-20
A	CAPECCHI, M.R. Targeted Gene Replacment. Scientific American. March 1994. pages 34-41.	1-20
A	MOREADITH et al. Gene Targeting in Embryonic Stem Cells: The New Physiology and Metabolism. J. Mol. Med. 1997. Vol. 75. pages 208-216.	1-20

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US01/09855

## A. CLASSIFICATION OF SUBJECT MATTER: IPC (7):

A01N 63/00, 65/00; A61K 48/00; G01N 33/00; A01K 67/00, 67/027, 67/033; C12N 15/00

## B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

WEST 2.0, SCISEARCH, JAPIO, MEDLINE, EMBASE, CAPLUS, BIOSIS, USPATFULL  
search terms; PDGF-C, transgenic non-human animal, PDGFC, PDGF, eriksson, aase, ponten, overexpression,  
martin, endothelial cells, platelets, fish oils, protease-activated ligand, platelet derived growth factor, thrombin



**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**

**THIS PAGE BLANK (USPTO)**